

Fear conditioning biases olfactory stem cell receptor fate

Clara W. Liff, Yasmine R. Ayman, Eliza C.B. Jaeger, Hudson S. Lee, Alexis Kim, Angélica Viña Albarracín, Bianca Jones Marlin 

Mortimer B. Zuckerman Mind Brain and Behavior Institute, Columbia University; New York, 10027, USA • Howard Hughes Medical Institute, Columbia University, New York, NY, USA • Department of Psychology, Columbia University; New York, 10032, USA

Reviewed Preprint

Published from the original preprint after peer review and assessment by eLife.

About eLife's process

Reviewed preprint posted

December 15, 2023 (this version)

Sent for peer review

September 28, 2023

Posted to bioRxiv

September 19, 2023


 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

The main olfactory epithelium initiates the process of odor encoding. Recent studies have demonstrated intergenerationally inherited changes in the olfactory system in response to fear conditioning, resulting in increases in olfactory receptor frequencies and altered responses to odors. We investigated changes in the morphology of the olfactory epithelium in response to an aversive stimulus. Here, we achieve volumetric cellular resolution to demonstrate that olfactory fear conditioning increases the number of odor-encoding neurons in mice that experience odor-shock conditioning (F0), *as well as their offspring* (F1). We provide evidence that increases in F0 were due to biased stem cell receptor choice. Thus, we reveal dynamic regulation of the olfactory epithelium receptor composition in response to olfactory fear conditioning, providing insight into the heritability of acquired phenotypes.

Graphical Abstract



One-Sentence Summary

Odor-shock pairing is inherited by naïve offspring and biases neurogenesis in the nose.

Highlights

- Olfactory fear conditioning leads to an increase in conditioned-odor-responsive cells in parents (F0) that is heritable (F1)
- Increase in conditioned-odor-responsive cells is sustained through at least 9 weeks of cell turnover in the main olfactory epithelium
- Olfactory fear conditioning in F0 biases neurogenesis specifically toward conditioned-odor responsive cell fate

eLife assessment

This **important** study seeks to advance the current understanding of intergenerational olfactory changes associated with odor-induced fear conditioning in mice. Whilst the overall approach employed by the authors is appropriate and the evidence presented in support of claims is **solid**, there is general agreement that specific points - particularly the lack of effect in the F1 generation - deserve further attention.

Main Text

Aversive olfactory conditioning in mice results in the persistent avoidance of the conditioned odor, and the olfactory sensory neurons (OSNs) responsive to this odor increase in number in the sensory epithelium (1). Strikingly, this increase in the number of specific sensory neurons was observed not only in trained F0 males, but also in their offspring (F1), despite the fact that the progeny had never been exposed to the conditioned odor (2–4). This phenomenon, intergenerational epigenetic inheritance, invokes the transfer of information from one generation to the next without alterations to the sequence of the genome.

Transgenerational epigenetic inheritance, the transfer of information beyond the F1 generation, is responsible for several examples of non-Mendelian transmission in plants, fission yeast, and worms (5–9). Molecular genetics has provided a detailed mechanistic understanding of the transmission of epigenetic information from parent to offspring in these organisms (10, 11). Olfactory conditioning in the parent may provide future generations with an adaptive advantage: enhanced sensitivity to aversive sensory features in the environment of the parent. Intergenerational epigenetic inheritance of olfactory properties in mice poses an elusive problem as to how signals responsible for the increase in specific OSNs in the sensory epithelium are transmitted from the nose, to the gamete, and then to the offspring.

Olfactory perception is initiated by the recognition of odorants by a large repertoire of receptors in the sensory epithelium. Individual sensory neurons in mice express only one of 1,400 different receptor genes (12). The choice of a receptor is stochastic and is mediated by an unusual mechanism of transactivation that delivers the necessary transcription factors to only one allele of a single receptor gene in a sensory neuron (13–15). Neurons expressing a given receptor are distributed within zones of the epithelium but project with precision to spatially invariant glomeruli in the olfactory bulb. Each odorant can interact with multiple distinct receptors, resulting in the activation of a unique ensemble of glomeruli. The recognition of an odor requires the integration of information from multiple glomeruli to the mitral and tufted cells in the olfactory bulb, and then to downstream olfactory convergent areas (16–19). If the stochastic choice of a single receptor in each neuron can be biased by salient odor associations in the environment, this would afford a mechanism to alter receptor representations in the epithelium.

The mechanisms responsible for the increase in the number of OSNs following aversive conditioning are more readily addressed in the sensory epithelium of F0 mice than in the F1 progeny. Elucidation of these local signaling events may then provide insight into the more distant transmission of information to the gametes. The mature olfactory sensory epithelium undergoes constant neurogenesis throughout the life of vertebrates. In mice, the lifespan of a mature OSN is estimated to be 30 days, and new sensory neurons are continually generated by the division of

basal stem cells, transit amplification, and ultimately differentiation to the mature OSN (20). Continual balanced neurogenesis suggests that increases in OSNs upon aversive conditioning could result from the increased birth or enhanced survival of a specific OSN population.

In this study, we performed quantification of OSNs after aversive olfactory learning, corroborating the results of an increase in the OSNs expressing a receptor responsive to the conditioned odor. Moreover, enhanced numbers of these specific OSNs are also observed in the F1 generation. Contrary to previous studies, we do not observe the inheritance of odor-evoked aversion to the conditioned odor in the F1 generation using our behavioral paradigm. We demonstrate that this phenomenon is persistent, as increases in specific OSNs continue for at least 63 days after conditioning. We further demonstrate that a biased increase in specific OSNs after learning is likely to result from the enhanced birth of specific OSNs, suggesting that biased receptor choice underlies this phenomenon in the parent and is epigenetically inherited by their offspring.

Olfactory fear conditioning leads to an increase in conditioned-odor-responsive cells in parents (F0)

In initial experiments, we asked whether we could observe changes in the abundance of receptors responsive to conditioned odors after aversive olfactory learning. The odorant receptor M71 is responsive to the odorant acetophenone, whereas neurons expressing MOR23 are activated by lyral. Homozygous mice modified at the M71 or MOR23 loci to also express GFP allowed for a determination of receptor abundance. These mice were subjected to an aversive olfactory conditioning paradigm in which acetophenone, lyral, or propanol, co-terminating with 0.75mA foot shock, were presented 5 times daily for 3 consecutive days (Fig. 1B,C). The unpaired control group also received odor presentations but experienced a 60-second delay prior to foot shock (Fig. 1C). Only mice in which odorant and shock were paired exhibited conditioned aversive behavior (Fig. S1B,C,D. Tukey's multiple comparisons. F0 acetophenone unpaired vs. paired $P < 0.0001$. $n = 10, 15$. F0 lyral unpaired vs. paired $P < 0.0001$. $n = 17, 20$. F0 propanol unpaired vs. paired $P < 0.0001$. $n = 16, 20$). Mouse nasal turbinates were surgically extracted 21 days after the initiation of training and subjected to iDISCO+ tissue clearing to visualize M71- and MOR23-expressing OSNs in transparent intact olfactory epithelia (Fig. 1B,D,E). We then imaged the cleared epithelia using light sheet microscopy and counted the number of M71 or MOR23 OSNs in a fixed volume of tissue using automated spot detection software (Fig. 1G,I).

Importantly, both M71 and MOR23 OSNs are expressed in the same zone of the epithelium, enabling consistent imaging and counting protocols for both OSN populations. Male and female M71-IRES-tauGFP^{+/+} (M71GFP) mice paired with acetophenone exhibited a 33% increase in the number of M71 OSNs 21 days after the initiation of aversive conditioning when compared to unpaired controls (Fig. 1H. One-way ANOVA. $P < 0.0001$. Tukey's multiple comparisons. Naïve vs. F0 paired $P < 0.0001$. F0 unpaired vs. F0 paired $P < 0.0001$. $n = 12, 11, 12$). Male and female MOR23-IRES-tauGFP^{+/+} (MOR23GFP) mice conditioned with lyral exhibited a 39% increase in MOR23 OSNs when compared to unpaired controls (Fig. 1J. One-way ANOVA. $P < 0.0001$. Tukey's multiple comparisons. Naïve vs. F0 paired $P < 0.001$. F0 unpaired vs. F0 paired $P < 0.0001$. $n = 7, 9, 9$). We performed a series of control experiments to demonstrate that fear conditioning does not lead to a global increase in the number of OSNs per cubic volume in the sensory epithelium. The odorant propanol does not activate M71 sensory neurons (1, 22). When propanol was employed as the conditioned odor in both the paired and unpaired training paradigms, we observed no difference in the number of M71 OSNs between the two groups of mice (Fig. 1F. Student's unpaired t-test. Unpaired vs. paired $P = 0.3009$. $n = 6, 7$). These results indicate that olfactory fear conditioning results in a specific increase in the number of cells responsive to the conditioned odor and does not lead to a global increase in all OSNs.




Fig. 1.

Olfactory fear conditioning leads to an increase in conditioned-odor-responsive cells in parents (F0) that is heritable (F1).

(A) Schematic representation of the mouse main olfactory epithelium and olfactory bulb. MOE: main olfactory epithelium. OB: olfactory bulb. **(B)** Timeline of olfactory fear conditioning and MOE collection. **(C)** Experimental paradigms for olfactory fear conditioning groups. Mice in the paired condition received a foot shock that co-terminated with odor presentation, while mice in the unpaired condition received a foot shock 60 seconds after odor presentation. **(D)** Schematic demonstrating the process by which cells of interest in the MOE were quantified. Epithelia from both M71-IRES-tauGFP^{+/+} and MOR23-IRES-tauGFP^{+/+} adult mice were cleared using the iDISCO+ tissue-clearing protocol. Samples were imaged on a light sheet microscope and analyzed using Imaris spot detection software. **(E)** Images of the MOE before (left) and after (right) optical tissue clearing. **(F)** The average number of M71 olfactory sensory neurons in a 350³ μm³ cube of the epithelium in the propanol unpaired (light blue) and propanol paired (dark blue) conditions (Student's unpaired t-test. Unpaired vs. paired P=0.3009. n=6,7.). **(G)** Example images of M71 OSNs in zone 1 of cleared MOE from both the unpaired (left) and paired (middle) conditions. Example image of an MOE with the counted cells represented by colored dots (right). Each set of colors represents a distinct counting cube. Scale bar: 200μm. **(H)** Graph showing the differences between the average number of M71 OSNs in a 350³ μm³ cube of epithelium of naive (gray), acetophenone unpaired (lighter green), and acetophenone paired (darker green) conditions in F0 and F1 (One-way ANOVA. P<0.0001. Tukey's multiple comparisons. Naive vs. F0 paired P<0.0001. F0 unpaired vs. paired P<0.0001. Naive vs. F1 paired P<0.0001. F1 unpaired vs. paired P<0.0001. n=12,11,12,12,14.). **(I)** Example images of MOR23 OSNs in zone 1 of cleared MOE from both the unpaired (left) and paired (middle) conditions. Example image of an MOE with the counted cells represented by colored dots (right). Scale bar: 200μm. **(J)** Graph showing the differences between the average number of MOR23 OSNs in a 350³ μm³ cube of epithelium in naive (gray), lyral unpaired (lighter purple), and lyral paired (darker purple) conditions in F0 and F1 (One-way ANOVA. P<0.0001. Tukey's multiple comparisons. Naive vs. F0 paired P<0.0001. F0 unpaired vs. paired P<0.0001. F1 unpaired vs. paired P=0.0368. n=7,9,9,6,6.).

Conditioned-odor-responsive cell increase is sustained through at least 9 weeks of cell turnover

We observe an increase in M71 OSNs 21 days after aversive conditioning with acetophenone. We next asked if this increase persists at later time points ([Fig. 2A](#)). At 42 days, we observe a persistent 20% increase ([Fig. 2B](#), Tukey's multiple comparisons, 42d unpaired vs. paired $P=0.0476$, $n=8,8$.) in the number of M71 OSNs in paired versus unpaired animals and a 30% increase at 63 days ([Fig. 2B](#), Tukey's multiple comparisons, 63d unpaired vs. paired $P=0.0011$, $n=4,6$.). Since the half-life ($t_{1/2}$) of the mouse olfactory sensory epithelium (the amount of time required for half of the epithelium to regenerate) is approximately 26 days ([23](#)), then at 42 days, approximately 67% will have been replaced by newly born neurons, and at 63 days, approximately 81% will have been replaced. The observation that these changes persist at least 63 days after aversive conditioning, together with the reported 26-day half-life for the main olfactory epithelium, suggests that a signaling mechanism must persist despite the fact that the entire sensory epithelium present at the time of conditioning will eventually be regenerated.

Fear conditioning-induced increases in conditioned-odor-responsive cells is heritable (F1)

We next asked whether the increase in the number of specific OSNs observed following conditioning is inherited by naïve offspring. Ten days after the initiation of aversive training, we bred F0 males from both the paired and unpaired groups with naïve M71GFP^{+/+} or MOR23GFP^{+/+} female mice. Each mating pair was separated ten days after co-housing to ensure that the offspring were never exposed to the conditioned father. Main olfactory epithelia were then collected from the offspring of these mating pairs at 8 weeks of age. The F1 mice were never exposed to acetophenone or lyral, nor had they undergone aversive conditioning. We nonetheless observed a 36% increase in M71 OSNs in both male and female offspring whose fathers experienced paired aversive conditioning with acetophenone when compared with the F1 of fathers that experienced the unpaired training paradigm ([Fig. 1H](#), Tukey's multiple comparisons, F1 unpaired vs. F1 paired $P<0.0001$, $n=12,14$.). A similar relative increase of 27% was observed in MOR23 OSNs in offspring of fathers that experienced paired aversive conditioning with lyral compared to F1 of unpaired fathers ([Fig. 1J](#), Tukey's multiple comparisons, F1 unpaired vs. F1 paired $P=0.0368$, $n=6,6$.). These results demonstrate the intergenerational epigenetic inheritance of an olfactory phenotype, namely an increase in specific OSNs in naïve F1 offspring following aversive conditioning in F0.

Previous behavioral studies demonstrated that offspring from fathers that experienced aversive olfactory conditioning exhibit enhanced sensitivity to the conditioned odor in both odor potentiated startle and aversive odor association assays ([2](#)). Therefore, we asked whether we could detect an aversive behavioral response to either acetophenone or lyral in the F1 population after aversive training in F0 fathers. In initial experiments, we performed aversive conditioning with either acetophenone or lyral in F0 males and females. Five days after the initiation of training, we placed mice in a 3-chamber arena with the conditioned odor on one side and a control odor (propanol) on the other. F0 mice in the paired group actively avoided the conditioned odor, whereas mice in the unpaired group exhibited no aversion to the conditioned odors ([Fig. S1B,C](#), Tukey's multiple comparisons, F0 acetophenone unpaired vs. paired $P=0.0008$, $n=10,15$, F0 lyral unpaired vs. paired $P=0.0057$, $n=10,15$.). Control mice spent roughly equal time exploring the propanol and conditioned odor chambers, whereas the paired mice spent approximately 67% (lyral paired) to 75% (acetophenone paired) of the time exploring the propanol chamber. Importantly, the offspring of F0 fathers that experienced aversive training with acetophenone exhibited no apparent avoidance of the conditioned odor. F1 mice spent equal time in both chambers ([Fig. S1B](#), Tukey's multiple comparisons, F1 acetophenone unpaired vs. paired $P=0.9551$, $n=4,6$.). We hypothesize that a higher number of conditioned odor-responsive OSNs in F1, despite no avoidance to the F0 paired odor, may position animals to learn avoidance behaviors in fewer




Fig. 2.

Conditioned-odor-responsive cell increase is sustained through at least 9 weeks of cell turnover.

(A) Timeline of olfactory fear conditioning and extended MOE collection time points. (B) The average number of M71 OSNs in a $350^3 \mu\text{m}^3$ cube of epithelium of unpaired (light green) and paired (dark green) mice, 42- or 63-days post-conditioning (One-way ANOVA. $P=0.0033$. Tukey's multiple comparisons. 42d unpaired vs. paired $P=0.0476$. 63d unpaired vs. paired $P=0.0011$. $n=8,8,4,6$).

trials or at lower odor concentrations. We note an unexplained result with the offspring of male mice conditioned with propanol. Propanol was behaviorally neutral in the F1 offspring from fathers that experienced paired aversive training with propanol (Fig. S1D. Tukey's multiple comparisons. Naïve vs. F1 paired $P=0.6624$. $n=25,12$.). However, offspring from fathers that had undergone unpaired training exhibited an attraction to propanol (Fig. S1D. Tukey's multiple comparisons. Naïve vs. F1 unpaired $P=0.0001$. $n=25,5$.). We note our avoidance paradigm may not be as sensitive to modest behavioral responses. Taken together, these results demonstrate that aversive odorant conditioning in the F0 population elicits active avoidance, but suggest that this behavioral phenotype is not transmitted to F1 progeny.

Olfactory fear conditioning biases olfactory receptor choice toward conditioned-odor-responsive cell-specific identities

The olfactory epithelium undergoes neurogenesis for the life of the organism. This continual renewal of OSNs suggests a possible mechanism for the observed increase in specific neuron populations responsive to conditioned odors. The increase in M71 and MOR23 cells following aversive training could result from a biased increase in either the birth or survival of specific OSNs. In initial experiments, we examined the relative number of M71 and MOR23 OSNs born during and after aversive training. We injected mice at the onset of training with 5-Ethynyl-2'-deoxyuridine (EdU), a thymidine analog that incorporates into newly synthesized DNA and labels newborn cells. Animals were injected during each of the 3 days of training and for 2 subsequent days (Fig. 3A). Epithelia were examined to determine the number of EdU-labeled M71 and MOR23 OSNs 21 days after the initiation of paired and unpaired aversive training (Fig. 3A). Since EdU has a half-life of approximately 35 minutes (24), analysis of EdU 16 days after the cessation of EdU exposure reflects a pulse-chase, allowing us to quantify a subset of the neurons born during the 5 days following the initiation of aversive conditioning (Fig. 3B).

The number of newborn M71 cells (EdU-labeled) out of total M71 cells is $1.24 \pm 0.29\%$ in naïve mice, $2.91 \pm 0.56\%$ after the unpaired paradigm, and $7.61 \pm 0.53\%$ following paired aversive training with acetophenone (Fig. 3D. One-way ANOVA. $P<0.0001$. Tukey's multiple comparisons. Naïve vs. paired $P<0.0001$. Unpaired vs. paired $P<0.0001$. $n=6,6,6$.). When lyral is used as the conditioned odor, the number of newborn MOR23 cells out of total MOR23 cells is $0.29 \pm 0.06\%$ in naïve mice, $0.55 \pm 0.09\%$ after the unpaired paradigm, and $1.11 \pm 0.21\%$ following paired aversive training (Fig. 3E. One-way ANOVA. $P=0.0120$. Tukey's multiple comparisons. Naïve vs. paired $P=0.0154$. Unpaired vs. paired $P=0.0653$. $n=4,6,8$.). These observations suggest that aversive learning results in a significant increase in the number of newborn M71 and MOR23 cells when comparing the paired and unpaired paradigms. The observation that the percentage of newborn M71 cells is 4-5 times that of MOR23 may simply reflect differences in the birth rates of the two cell populations.

We scored the number of EdU-labeled M71 and MOR23 OSNs 16 days after the cessation of EdU exposure. Since the differentiation of a newborn cell to a mature olfactory neuron requires about 7-10 days (20, 25), these data strongly suggest that aversive training results in a specific increase in the birth of new cells responsive to the conditioned odor. Experiments suggest that an enhanced rate of survival is not responsible for the observed increase in specific OSNs. If EdU is administered prior to the onset of training, an increase in the number of EdU+ cells would reflect enhanced survival rather than increased birth. However, daily exposure to EdU for 5 days 12 days prior to conditioning does not reveal a relative increase in the frequency of EdU+ M71 or MOR23 cells when comparing the paired and unpaired paradigms (data not shown). Taken together, these data strongly suggest that the specific increase in cells responsive to the conditioned odor is a consequence of a relative increase in the birth of new cells expressing the M71 and MOR23 receptors.




Fig. 3.

Olfactory fear conditioning biases olfactory receptor choice toward conditioned-odor-responsive cell-specific identities.

(A) Timeline of olfactory fear conditioning, EdU injections, and MOE collection. **(B)** Schematic representation of the distinct layers of the MOE, showing the stem cell, immature OSN, and mature OSN populations (left). Representative image of the MOE from an MOR23GFP^{+/+} mouse showing EdU-positive cells (red) and a newborn (EdU+) MOR23 OSN (green). Scale bar: 20 μ m. **(C)** Representative images showing staining of EdU (red, first column), endogenous GFP (green, second column), DAPI (blue, third column), and the merged channels (fourth column) in both M71GFP^{+/+} and MOR23GFP^{+/+} MOE. Scale bar: 40 μ m. **(D)** Percentage of EdU-positive M71 OSNs in naive, unpaired, and paired groups (One-way ANOVA. P<0.0001. Tukey's multiple comparisons. Naive vs. paired P<0.0001. Unpaired vs. paired P<0.0001. n=6,6,6.). **(E)** Percentage of EdU-positive MOR23 OSNs in naive, unpaired, and paired groups (One-way ANOVA. P=0.0120. Tukey's multiple comparisons. Naive vs. paired P=0.0154. Unpaired vs. paired P=0.0653. n=4,6,8.).

Discussion

We used tissue-clearing techniques and light-sheet microscopy to demonstrate an increase in olfactory sensory neurons (OSNs) expressing the receptor for an aversively conditioned odor. Moreover, enhanced cell numbers to the conditioned odor were observed in naïve offspring.

These data are in accord with findings that employ other cellular visualization techniques (2–4). In F0, this increase is stable for 63 days, a time by which the vast majority of the cells present during aversive training have been replaced by newborn sensory neurons. The increase in F0 results, in part, from the contribution of newborn neurons responsive to the conditioned odor, demonstrating a biasing of olfactory receptors during OSN development. The sustained increase in F0, along with the inheritance in F1, suggests that there is a stable signal that is responsible for the induction, maintenance, and inheritance of the increase in OSNs responsive to the paired odor.

The stochastic choice of olfactory receptors may provide an opportunity to alter the representation of receptors in order to allow an organism to adapt to the environment. Changes in the number of OSNs may lead to an increase in sensitivity of the paired odor. A change in OSN number may also lead to increased inputs to downstream sensory areas. Such perceptual changes have been reported in the motor, visual, olfactory, and auditory systems, where topographical arrangements at the primary sensory cortex are modulated in certain fear conditioning paradigms in mammals (26–31), although this has not yet been demonstrated intergenerationally. We observed an increase in conditioned odor-responsive neurons in both the F0 and F1 populations. Only the F0, however, exhibited avoidance behavior. We speculate that the increase in neurons responsive to the conditioned odor could enhance the sensitivity to, or the discrimination of, the paired odor in F0 and F1. This would enable the F1 population to learn that odor predicts shock with fewer training cycles or less odorant when trained with the conditioned odor. An area of exploration is how a long latency between training and testing (e.g., 42 or 63 days, timepoints at which we observe a sustained increase in cell number) affects avoidance behavior. These studies will lend insight to the behavioral differences observed in F1. These findings set a foundation to uncover the mechanism by which olfactory receptor bias is communicated within the main olfactory epithelium, to the germline, and, moreover, maintained during the development of offspring. What remains to be uncovered are the mechanisms to bias the choice of specific receptors in the main olfactory epithelium and how the information governing the biasing of receptor choice is transferred to the gametes.

In mice, the paternal transmission of epigenetic information has been observed following metabolic disturbances, social stress, and exposure to drugs and toxins (32). High-fat or low-protein diets, as well as caloric restriction in the father, results in metabolic disturbances in the offspring, even after *in vitro* fertilization (33, 34). Parental stressors, such as chronic defeat or maternal separation, result in hormonal disturbances and behavioral phenotypes in the offspring (35–37). Finally, toxins and addictive drugs result in an array of metabolic disturbances in the F1 population that recapitulate the paternal state (38). These paternal stressors are associated with metabolic and hormonal disturbances that can readily act at a distance to affect the gamete. It has been demonstrated in male gametogenesis that extracellular vesicles in the testes transmit an RNA payload as they fuse with maturing sperm (39–42). Such studies provide insights into a mechanism by which an olfactory sensory experience paired with fear learning could transmit receptor-specific information from one generation to the next.

Our study elaborates on a function of sensory systems in which a learned adaptation can influence future generations. Thus, the distinction between innate and learned behaviors may be fundamentally flexible — learned adaptations in the parent may have the potential to become innate in their offspring. Understanding the mechanisms of inherited adaptation will provide insight for interventions when these changes no longer serve as adaptive to the organism.

Acknowledgements

We thank the past and current members of the Marlin Lab for critical discussions and inputs on the manuscript. We thank Dr. Richard Axel, Dr. Stavros Lomvardas, Dr. Carol Mason, Dr. Robert Froemke, Dr. Ishmail Abdus-Saboor, Dr. Ismail Ahmed, and Dr. Joseph Jones Marlin for their comments, discussion, and technical assistance. We thank A.I.S.C, A.M., and V.L.F.N. for their support. We thank Stellate Communications for assistance with figure production and preparation of the manuscript. Imaging was performed with support from the Zuckerman Institute's Cellular Imaging platform.

Funding

Simons Society of Fellows Junior Fellowship 524991(BJM)

UNCF E.E. Just Fellowship CU20-1071 (BJM)

BBRF NARSAD Young Investigator Grant 30380 (BJM)

National Institutes of Health grant 1S10OD023587-01

Author contributions

Conceptualization: CWL, YRA, BJM

Methodology: CWL, YRA, ECBJ, HSL,

BJM Investigation: CWL, YRA, ECBJ, HSL, AK, BJM

Visualization: CWL, YRA

Funding acquisition: BJM

Project administration: CWL, BJM

Supervision: BJM

Writing – original draft: CWL, YRA, BJM

Writing – review & editing: CWL, AVA, BJM

Competing interests

Authors declare that they have no competing interests.

Data and materials availability

All data are available upon request.

Supplementary Materials

Materials and Methods

Figs. S1 to S7

Movies S1 to S8

References

1. Jones S. V., Choi D. C., Davis M., Ressler K. J. (2008) **Learning-Dependent Structural Plasticity in the Adult Olfactory Pathway** *J Neurosci* **28**:13106–13111
2. Dias B. G., Ressler K. J. (2014) **Parental olfactory experience influences behavior and neural structure in subsequent generations** *Nat Neurosci* **17**:89–96
3. Aoued H. S. *et al.* (2019) **Reversing Behavioral, Neuroanatomical, and Germline Influences of Intergenerational Stress** *Biol. Psychiatry* **85**:248–256
4. Aoued H. S., Sannigrahi S., Hunter S. C., Doshi N., Sathi Z. S., Chan A. W. S., Walum H., Dias B. G. (2020) **Proximate causes and consequences of intergenerational influences of salient sensory experience** *Genes, Brain and Behavior* **19**
5. Greer E. L., Maures T. J., Ucar D., Hauswirth A. G., Mancini E., Lim J. P., Benayoun B. A., Shi Y., Brunet A. (2011) **Transgenerational epigenetic inheritance of longevity in *Caenorhabditis elegans*** *Nature* **479**:365–371
6. Schmitz R. J., Schultz M. D., Lewsey M. G., O'Malley R. C., Urich M. A., Libiger O., Schork N. J., Ecker J. R. (2011) **Transgenerational Epigenetic Instability Is a Source of Novel Methylation Variants** *Science* **334**:369–373
7. Rechavi O., Houry-Ze'evi L., Anava S., Goh W. S. S., Kerk S. Y., Hannon G. J., Hobert O. (2014) **Starvation-Induced Transgenerational Inheritance of Small RNAs in *C. elegans*** *Cell* **158**:277–287
8. Yu R., Wang X., Moazed D. (2018) **Epigenetic inheritance mediated by coupling of RNAi and histone H3K9 methylation** *Nature* **558**:615–619
9. Moore R. S., Kaletsky R., Lesnik C., Cota V., Blackman E., Parsons L. R., Gitai Z., Murphy C. T. (2021) **The role of the Cer1 transposon in horizontal transfer of transgenerational memory** *Cell* **184**:4697–4712
10. Rando O. J., Verstrepen K. J. (2007) **Timescales of genetic and epigenetic inheritance** *Cell* **128**:655–668
11. Fitz-James M. H., Cavalli G. (2022) **Molecular mechanisms of transgenerational epigenetic inheritance** *Nat Rev Genet* **23**:325–341
12. Buck L., Axel R. (1991) **A novel multigene family may encode odorant receptors: a molecular basis for odor recognition** *Cell* **65**:175–187
13. Chess A., Simon I., Cedar H., Axel R. (1994) **Allelic inactivation regulates olfactory receptor gene expression** *Cell* **78**:823–834
14. Shykind B. M., Rohani S. C., O'Donnell S., Nemes A., Mendelsohn M., Sun Y., Axel R., Barnea G. (2004) **Gene Switching and the Stability of Odorant Receptor Gene Choice** *Cell* **117**:801–815
15. Lomvardas S., Barnea G., Pisapia D. J., Mendelsohn M., Kirkland J., Axel R. (2006) **Interchromosomal Interactions and Olfactory Receptor Choice** *Cell* **126**:403–413

16. Price J. L., Powell T. P. S. (1970) **The Mitral and Short Axon Cells of the Olfactory Bulb** *Journal of Cell Science* **7**:631–651
17. Price J. L. (1985) **Beyond the primary olfactory cortex: olfactory-related areas in the neocortex, thalamus and hypothalamus** *Chemical Senses* **10**:239–258
18. Chen C.-F. F., Zou D.-J., Altomare C. G., Xu L., Greer C. A., Firestein S. J. (2014) **Nonsensory target-dependent organization of piriform cortex** *Proceedings of the National Academy of Sciences* **111**:16931–16936
19. Diodato A., De Brimont M., Ruinart, Yim Y. S., Derian N., Perrin S., Pouch J., Klatzmann D., Garel S., Choi G. B., Fleischmann A. (2016) **Molecular signatures of neural connectivity in the olfactory cortex** *Nature Communications* **7** <https://doi.org/10.1038/ncomms12238>
20. Liberia T., Martin-Lopez E., Meller S. J., Greer C. A. **T. Liberia, E. Martin-Lopez, S. J. Meller, C. A. Greer, eNeuro, in press, doi:10.1523/ENEURO.0266-19.2019.** <https://doi.org/10.1523/ENEURO.0266-19.2019>
21. Renier N. *et al.* (2016) **Mapping of brain activity by automated volume analysis of immediate early genes** *Cell* **165**:1789–1802
22. Johnson B. A., Leon M. (2000) **Modular representations of odorants in the glomerular layer of the rat olfactory bulb and the effects of stimulus concentration** *Journal of Comparative Neurology* **422**:496–509
23. Holl A.-M. (2018) **Survival of mature mouse olfactory sensory neurons labeled genetically perinatally** *Mol Cell Neurosci* **88**:258–269
24. Cheraghali A. M., Kumar R., Knaus E. E., Wiebe L. I. (1995) **Pharmacokinetics and bioavailability of 5-ethyl-2'-deoxyuridine and its novel (5R,6R)-5-bromo-6-ethoxy-5,6-dihydro prodrugs in mice** *Drug Metab Dispos* **23**:223–226
25. Miragall F., Graziadei G. A. Monti (1982) **Experimental studies on the olfactory marker protein. II. Appearance of the olfactory marker protein during differentiation of the olfactory sensory neurons of mouse: an immunohistochemical and autoradiographic study** *Brain Res* **239**:245–250
26. Ressler K. J., Sullivan S. L., Buck L. B. (1994) **Information coding in the olfactory system: Evidence for a stereotyped and highly organized epitope map in the olfactory bulb** *Cell* **79**:1245–1255
27. Vassar R., Chao S. K., Sitcheran R., Nun-ez J. M., Vosshall L. B., Axel R. (1994) **Topographic organization of sensory projections to the olfactory bulb** *Cell* **79**:981–991
28. Mombaerts P., Wang F., Dulac C., Chao S. K., Nemes A., Mendelsohn M., Edmondson J., Axel R. (1996) **Visualizing an Olfactory Sensory Map** *Cell* **87**:675–686
29. Lai C. S. W., Adler A., Gan W.-B. (2018) **Fear extinction reverses dendritic spine formation induced by fear conditioning in the mouse auditory cortex** *Proceedings of the National Academy of Sciences* **115**:9306–9311
30. Xu Z., Adler A., Li H., Pérez-Cuesta L. M., Lai B., Li W., Gan W.-B. (2019) **Fear conditioning and extinction induce opposing changes in dendritic spine remodeling and somatic activity of layer 5 pyramidal neurons in the mouse motor cortex** *Sci Rep* **9**

31. Li Z., Yan A., Guo K., Li W. (2019) **Fear-Related Signals in the Primary Visual Cortex** *Current Biology* **29**:4078–4083
32. Huypens P., Sass S., Wu M., Dyckhoff D., Tschöp M., Theis F., Marschall S., de Angelis M. Hrabě, Beckers J. (2016) **Epigenetic germline inheritance of diet-induced obesity and insulin resistance** *Nat Genet* **48**:497–499
33. Carone B. R. *et al.* (2010) **Paternally induced transgenerational environmental reprogramming of metabolic gene expression in mammals** *Cell* **143**:1084–1096
34. Chen Q. *et al.* (2016) **Sperm tsRNAs contribute to intergenerational inheritance of an acquired metabolic disorder** *Science* **351**:397–400
35. Dietz D. M., Laplant Q., Watts E. L., Hodes G. E., Russo S. J., Feng J., Oosting R. S., Vialou V., Nestler E. J. (2011) **Paternal transmission of stress-induced pathologies** *Biol Psychiatry* **70**:408–414
36. Morgan C. P., Bale T. L. (2011) **Early Prenatal Stress Epigenetically Programs Dymasculinization in Second-Generation Offspring via the Paternal Lineage** *J. Neurosci* **31**:11748–11755
37. Gapp K., Jawaid A., Sarkies P., Bohacek J., Pelczar P., Prados J., Farinelli L., Miska E., Mansuy I. M. (2014) **Implication of sperm RNAs in transgenerational inheritance of the effects of early trauma in mice** *Nat Neurosci* **17**:667–669
38. Toussaint A. B. *et al.* (2022) **Chronic paternal morphine exposure increases sensitivity to morphine-derived pain relief in male progeny** *Sci Adv* **8**
39. Rando O. J. (2016) **Intergenerational Transfer of Epigenetic Information in Sperm** *Cold Spring Harb Perspect Med* **6**
40. Sharma U., Sun F., Conine C. C., Reichholf B., Kukreja S., Herzog V. A., Ameres S. L., Rando O. J. (2018) **Small RNAs Are Trafficked from the Epididymis to Developing Mammalian Sperm** *Dev Cell* **46**:481–494
41. Morgan C. P., Chan J., Bale T. L. (2019) **Driving the next generation: Paternal lifetime experiences transmitted via extracellular vesicles and their small RNA cargo** *Biol Psychiatry* **85**:164–171
42. Chan J. C. *et al.* (2020) **Reproductive tract extracellular vesicles are sufficient to transmit intergenerational stress and program neurodevelopment** *Nat Commun* **11**

Article and author information

Clara W. Liff

Mortimer B. Zuckerman Mind Brain and Behavior Institute, Columbia University; New York, 10027, USA

Yasmine R. Ayman

Mortimer B. Zuckerman Mind Brain and Behavior Institute, Columbia University; New York, 10027, USA

Eliza C.B. Jaeger

Mortimer B. Zuckerman Mind Brain and Behavior Institute, Columbia University; New York, 10027, USA

Hudson S. Lee

Mortimer B. Zuckerman Mind Brain and Behavior Institute, Columbia University; New York, 10027, USA

Alexis Kim

Mortimer B. Zuckerman Mind Brain and Behavior Institute, Columbia University; New York, 10027, USA, Howard Hughes Medical Institute, Columbia University, New York, NY, USA

Angélica Viña Albarracín

Mortimer B. Zuckerman Mind Brain and Behavior Institute, Columbia University; New York, 10027, USA

Bianca Jones Marlin

Mortimer B. Zuckerman Mind Brain and Behavior Institute, Columbia University; New York, 10027, USA, Howard Hughes Medical Institute, Columbia University, New York, NY, USA, Department of Psychology, Columbia University; New York, 10032, USA

For correspondence: bjm2174@columbia.edu

Copyright

© 2023, Liff et al.

This article is distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editors

Reviewing Editor

Mario Penzo

National Institute of Mental Health, United States of America

Senior Editor

Kate Wassum

University of California, Los Angeles, United States of America

Reviewer #1 (Public Review):

Summary:

The study by Liff et al significantly advances our understanding of transgenerational olfactory changes resulting from fear conditioning, particularly in revealing elevated odor-encoding neurons in both conditioned mice (F0) and their progeny (F1). The authors attribute F0 increases to biased stem cell receptor selection, building upon the seminal work of Dias and Ressler (2014). While the dedication and use of novel histological techniques add strength to the study, there are notable weaknesses, including the need for clarification on discrepancies with previous findings, the decision to modify paradigms, and the presentation of behavioral data in supplementary materials.

Overall, the manuscript has strong potential but would benefit from addressing these weaknesses and minor recommendations to enhance its quality and contribution to the field.

Strengths:

- Significant contribution to understanding transgenerational olfactory changes induced by fear conditioning.
- Use of novel histological techniques and exploration of stem cell involvement adds depth to the study.

Weaknesses:

Discrepancies with previous findings need clarification, especially regarding the absence of similar behavioral effects in F1. Lack of discussion on the decision to modify paradigms instead of using the same model. Presentation of behavioral data in supplementary materials, with a recommendation to include behavioral quantification in main figures. Absence of quantification for freezing behavior, a crucial measure in fear conditioning.

Reviewer #2 (Public Review):

Summary:

The authors examined inherited changes to the olfactory epithelium produced by odor-shock pairings. The manuscript demonstrates that odor fear-conditioning biases olfactory bulb neurogenesis toward more production of the olfactory sensory neurons engaged by the odor-shock pairing. Further, the manuscript reveals that this bias remains in first-generation male and female progeny produced by trained parents. Surprisingly, there was a disconnect between the increased morphology of the olfactory epithelium for the conditioned odor and the response to odor presentation. The expectation based on previous literature and the morphological results was that F1 progeny would also show an aversion to the odor stimulus. However, the authors found that F1 progeny were not more sensitive to the odor compared to littermate controls.

Strengths:

The manuscript includes conceptual innovation and some technical innovation. The results validate previous findings that were deemed controversial in the field, which is a major strength of the work. Moreover, these studies were conducted using a combination of genetically modified animals and state-of-the-art imaging techniques, highlighting the rigorous nature of the research. Lastly, the authors provide novel mechanistic details regarding the remodeling of the olfactory epithelium, demonstrating that biased neurogenesis, as opposed to changes in survival rates, account for the increase in odorant receptors after training.

Weaknesses:

The main weakness is the disconnect between the morphological changes reported and the lack of change in aversion to the odorant in F1 progeny. The authors also do not address the mechanisms underlying the inheritance of the phenotype, which may lie outside of the scope of the present study.

Reviewer #3 (Public Review):

In their paper entitled "Fear conditioning biases olfactory stem cell receptor fate" Liff et al. address the still enigmatic (and quite fascinating) phenomenon of intergenerationally inherited changes in the olfactory system in response to odor-dependent fear conditioning.

In the abstract / summary, the authors raise expectations that are not supported by the data. For example, it is claimed that "increases in F0 were due to biased stem cell receptor choice." While an active field of study that has seen remarkable progress in the past decade, olfactory

receptor gene choice and its relevant timing in particular is still unresolved. Here, Liff et al., do not pinpoint at what stage during differentiation the "biased choice" is made.

Similarly, the concluding statement that the study provides "insight into the heritability of acquired phenotypes" is somewhat misleading. The experiments do not address the mechanisms underlying heritability.

The statement that "the percentage of newborn M71 cells is 4-5 times that of MOR23 may simply reflect differences in the birth rates of the two cell populations" should, if true, result in similar differences in the occurrence of mature OSNs with either receptor identity. According to Fig. 1H & J, however, this is not the case.

An important result is that Liff et al., in contrast to results from other studies, "do not observe the inheritance of odor-evoked aversion to the conditioned odor in the F1 generation." This discrepancy needs to be discussed.

The authors speculate that "the increase in neurons responsive to the conditioned odor could enhance the sensitivity to, or the discrimination of, the paired odor in F0 and F1. This would enable the F1 population to learn that odor predicts shock with fewer training cycles or less odorant when trained with the conditioned odor." This is a fascinating idea that, in fact, could have been readily tested by Liff and coworkers. If this hypothesis were found true, this would substantially enhance the impact of the study for the field.